

TABLE 1.—FINAL 2004 ABCs, TACs, AND OVERFISHING LEVELS OF GROUND FISH FOR THE WESTERN/CENTRAL/WEST YAKUTAT (W/C/WYK), WESTERN (W), CENTRAL (C), EASTERN (E) REGULATORY AREAS, AND IN THE WEST YAKUTAT (WYK), SOUTHEAST OUTSIDE (SEO), AND GULFWIDE (GW) DISTRICTS OF THE GULF OF ALASKA

[Values are in metric tons]

Species	Area <sup>1</sup>	ABC	TAC	Overfishing
Pollock <sup>2</sup> Shumagin .....	(610)	22,930	22,930	.....

TABLE 1.—FINAL 2004 ABCs, TACs, AND OVERFISHING LEVELS OF GROUND FISH FOR THE WESTERN/CENTRAL/WEST YAKUTAT (W/C/WYK), WESTERN (W), CENTRAL (C), EASTERN (E) REGULATORY AREAS, AND IN THE WEST YAKUTAT (WYK), SOUTHEAST OUTSIDE (SEO), AND GULFWIDE (GW) DISTRICTS OF THE GULF OF ALASKA—Continued

[Values are in metric tons]

Species	Area <sup>1</sup>	ABC	TAC	Overfishing
Chirikof .....	(620)	26,490	26,490	.....
Kodiak .....	(630)	14,040	14,040	.....
WYK .....	(640)	1,280	1,280	.....
Subtotal .....	W/C/ WYK	64,740	64,740	91,060
SEO .....	(650)	6,520	6,520	8,690
Total .....		71,260	71,260	99,750
Pacific cod <sup>3</sup> .....	W	22,610	16,957	.....
	C	35,800	27,116	.....
	E	4,400	3,960	.....
Total .....		62,810	48,033	102,000
Flatfish <sup>4</sup> (deep-water) .....	W	310	310	.....
	C	2,970	2,970	.....
	WYK	1,880	1,880	.....
	SEO	910	910	.....
Total .....		6,070	6,070	8,010
Rex sole .....	W	1,680	1,680	.....
	C	7,340	7,340	.....
	WYK	1,340	1,340	.....
	SEO	2,290	2,290	.....
Total .....		12,650	12,650	16,480
Flathead sole .....	W	13,410	2,000	.....
	C	34,430	5,000	.....
	WYK	3,430	3,430	.....
	SEO	450	450	.....
Total .....		51,270	10,880	64,750
Flatfish <sup>5</sup> (shallow-water) .....	W	21,580	4,500	.....
	C	27,250	13,000	.....
	WYK	2,030	2,030	.....
	SEO	1,210	1,210	.....
Total .....		52,070	20,740	63,840
Arrowtooth flounder .....	W	23,590	8,000	.....
	C	151,840	25,000	.....
	WYK	10,590	2,500	.....
	SEO	8,910	2,500	.....
Total .....		194,930	38,000	228,130
Sablefish <sup>6</sup> .....	W	2,930	2,930	.....
	C	7,300	7,300	.....
	WYK	2,550	2,550	.....
	SEO	3,770	3,770	.....
Subtotal .....	E	6,320	6,320	.....
Total .....		16,550	16,550	22,160
Pacific ocean perch <sup>7</sup> .....	W	2,520	2,520	2,990
	C	8,390	8,390	9,960
	WYK	830	830	.....
	SEO	1,600	1,600	.....
Subtotal .....	E	.....	.....	2,890
Total .....		13,340	13,340	15,840
Shortraker/rougheye <sup>8</sup> .....	W	254	254	.....
	C	656	656	.....
	E	408	408	.....
Total .....		1,318	1,318	2,510
Other rockfish <sup>9 10</sup> .....	W	40	40	.....
	C	300	300	.....
	WYK	130	130	.....
	SEO	3,430	200	.....
Total .....		3,900	670	5,150

TABLE 1.—FINAL 2004 ABCS, TACS, AND OVERFISHING LEVELS OF GROUND FISH FOR THE WESTERN/CENTRAL/WEST YAKUTAT (W/C/WYK), WESTERN (W), CENTRAL (C), EASTERN (E) REGULATORY AREAS, AND IN THE WEST YAKUTAT (WYK), SOUTHEAST OUTSIDE (SEO), AND GULFWIDE (GW) DISTRICTS OF THE GULF OF ALASKA—Continued

[Values are in metric tons]

Species	Area <sup>1</sup>	ABC	TAC	Overfishing
Northern Rockfish <sup>10 12 15</sup> .....	W	770	770	.....
	C	4,100	4,100	.....
	E	N/A	N/A	.....
Total .....		4,870	4,870	5,790
Pelagic shelf rockfish <sup>13</sup> .....	W	370	370	.....
	C	3,010	3,010	.....
	WYK	210	210	.....
	SEO	880	880	.....
Total .....		4,470	4,470	5,570
Thornyhead rockfish .....	W	410	410	.....
	C	1,010	1,010	.....
	E	520	520	.....
Total .....		1,940	1,940	2,590
Demersal shelf rockfish <sup>11</sup> .....	SEO	450	450	690
Atka mackerel .....	GW	600	600	6,200
Other species <sup>14</sup> .....	GW	N/A	12,592	N/A
Total <sup>16</sup> .....		498,948	264,433	649,460

<sup>1</sup> Regulatory areas and districts are defined at § 679.2.

<sup>2</sup> Pollock is apportioned in the Western and Central Regulatory Areas among three statistical areas. During the A season, the apportionment is based upon an adjusted estimate of the relative distribution of pollock biomass at 23.62 percent, 56.9 percent, and 19.48 percent in Statistical Areas 610, 620, and 630, respectively. During the B season, the apportionment is based on the relative distribution of pollock biomass at 23.62 percent, 64.47 percent, and 8.91 percent in Statistical Areas 610, 620, and 630, respectively. During the C and D seasons, pollock is apportioned based on the relative distribution of pollock biomass at 48.64 percent, 21.3 percent, and 30.6 percent in Statistical Areas 610, 620, and 630, respectively. These seasonal apportionments are shown in Table 3. In the West Yakutat and Southeast Outside Districts of the Eastern Regulatory Area, pollock is not divided into seasonal allowances.

<sup>3</sup> The annual Pacific cod TAC is apportioned 60 percent to the A season and 40 percent to the B season in the Western and Central Regulatory Areas of the GOA. Pacific cod is allocated 90 percent for processing by the inshore component and 10 percent for processing by the offshore component. Seasonal apportionments and component allocations of TAC are shown in Table 4.

<sup>4</sup> “Deep water flatfish” means Dover sole, Greenland turbot, and deepsea sole.

<sup>5</sup> “Shallow water flatfish” means flatfish not including “deep water flatfish”, flathead sole, rex sole, or arrowtooth flounder.

<sup>6</sup> Sablefish is allocated to trawl and hook-and-line gears (Table 2).

<sup>7</sup> “Pacific ocean perch” means *Sebastes alutus*.

<sup>8</sup> “Shortraker/rougheye rockfish” means *Sebastes borealis* (shortraker) and *S. aleutianus* (rougheye).

<sup>9</sup> “Other rockfish” in the Western and Central Regulatory Areas and in the West Yakutat District means slope rockfish and demersal shelf rockfish. The category “other rockfish” in the Southeast Outside District means slope rockfish.

<sup>10</sup> “Slope rockfish” means *Sebastes aurora* (aurora), *S. melanostomus* (blackgill), *S. paucispinis* (bocaccio), *S. goodei* (chilipepper), *S. crameri* (darkblotch), *S. elongatus* (greenstriped), *S. variegatus* (harlequin), *S. wilsoni* (pygmy), *S. babcocki* (redbanded), *S. proriger* (redstripe), *S. zacentrus* (sharpchin), *S. jordani* (shortbelly), *S. brevispinis* (silvergrey), *S. diploproa* (splitnose), *S. saxicola* (stripetail), *S. miniatus* (vermillion), and *S. reedi* (yellowmouth). In the Eastern GOA only, “slope rockfish” also includes northern rockfish, *S. polyspinus*.

<sup>11</sup> “Demersal shelf rockfish” means *Sebastes pinniger* (canary), *S. nebulosus* (china), *S. caurinus* (copper), *S. maliger* (quillback), *S. helvomaculatus* (rosethorn), *S. nigrocinctus* (tiger), and *S. ruberrimus* (yelloweye).

<sup>12</sup> “Northern rockfish” means *Sebastes polyspinus*.

<sup>13</sup> “Pelagic shelf rockfish” means *Sebastes ciliatus* (dusky), *S. entomelas* (widow), and *S. flavidus* (yellowtail).

<sup>14</sup> “Other species” means sculpins, sharks, skates, squid, and octopus. The TAC for “other species” equals 5 percent of the TACs of assessed target species.

<sup>15</sup> N/A means not applicable.

<sup>16</sup> The total ABC and OFL is the sum of the ABCs and OFLs for assessed target species.